

THERMODYNAMIC CHARACTERISTICS OF THE HO<sub>2</sub>  
RADICAL IN AN AQUEOUS SOLUTION

V.M. Berdnikov, O.S. Zhuraleva


Translation of: "Termodinamicheskiye Kharakteristiki radikala HO<sub>2</sub> v vodnom rastvore,"  
In: Zhurnal Fizicheskoy Khimii, v. 46, no. 10,  
1972, pp. 2658-61.

(NASA-TT-F-15793) THERMODYNAMIC  
CHARACTERISTICS OF THE HO<sub>2</sub> RADICAL IN AN  
AQUEOUS SOLUTION (Linguistic Systems,  
Inc., Cambridge, Mass.) 20-p HC \$4.00

N74-32579

Unclass

CSCL 07D G3/06 48436



NATIONAL AERONAUTICS AND SPACE ADMINISTRATION  
WASHINGTON, D.C. 20546 AUGUST 1974

1. Report No. NASA TT F-15793	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle THERMODYNAMIC CHARACTERISTICS OF THE HO <sub>2</sub> RADICAL IN AN AQUEOUS SOLUTION		5. Report Date August 1974	
		6. Performing Organization Code	
7. Author(s) V.M. Berdnikov, O.S. Zhuraleva		8. Performing Organization Report No.	
		10. Work Unit No.	
9. Performing Organization Name and Address Linguistic Systems Inc. Cambridge, MASS		11. Contract or Grant No. NASW-2482	
		13. Type of Report and Period Covered Translation	
12. Sponsoring Agency Name and Address NATIONAL AERONAUTICS & SPACE ADMINISTRATION WASHINGTON, D.C. 20546		14. Sponsoring Agency Code	
15. Supplementary Notes Translation of: "Termodinamicheskiye Kharakteristiki radikala HO <sub>2</sub> v vodnom rastvore," In: Zhurnal Fizicheskoy Khimii, v. 46, no. 10, 1972, pp. 2658-61.			
16. Abstract  An evaluation of the thermodynamic characteristics of the hydration and dissociation of HO <sub>2</sub> is presented.			
17. Key Words (Selected by Author(s))		18. Distribution Statement  Unlimited	
19. Security Classif. (of this report)	20. Security Classif. (of this page)	21. No. of Pages	22. Price


# THERMODYNAMIC CHARACTERISTICS OF THE HO<sub>2</sub> RADICAL IN AN AQUEOUS SOLUTION

V.M. Berdnikov, O.S. Zhuraleva

The thermodynamic characteristics of the hydration and dissociation of HO<sub>2</sub> were determined.

/2658\*

It was shown in [1] that the formation of radicals in the system Fe<sup>3+</sup> + H<sub>2</sub>O<sub>2</sub> occurs by the mechanism



Moreover the first two phases can be calculated by the equilibria


/2659

$$k_{\text{eff}} = \frac{k_1}{k_{-1}} \cdot \frac{k_2}{k_{-2}} \cdot k_3$$

The rate constant  $k_{\text{eff}} = 2 \cdot 10^{-5} \text{ sec}^{-1}$  at 25°C and at neutral ionic strength. In [2] it was found that the rate constant of the reaction  $\text{Fe}^{2+} + \text{HO}_2$  is equal to  $2.1 \cdot 10^6 / \text{mol} \cdot \text{sec}$  at 25°C and does not depend on the acidity of the medium, and the initial product of the reaction is the complex  $\text{Fe}^{3+} + \text{HO}_2^-$ . This indicates that the limiting phase here is the replacement of water in the coordination shell of the Fe<sup>2+</sup> ion. Thus the ratio can be written

$$K_{\text{e}} = \frac{k_1}{k_{-1}} \cdot \frac{k_2}{k_{-2}} \cdot \frac{k_3}{k_{-3}} = \frac{k_{\text{eff}}}{k_{-3}} = e^{-\Delta Z_0 / RT}$$

where  $K_p$  is the equilibrium constant of the reaction



\*Numbers in the margin indicate pagination in the foreign text.

We find  $K_p = 9.5 \cdot 10^{-12}$  and  $\Delta Z_O = 15.1$  kcal.

Knowing the standard redox potential of  $Fe^{3+}/Fe^{2+}$  (0.771 in [3]), we find the standard free energy  $\Delta Z_O = -32.9$  kcal for the reaction





Since  $\Delta Z_O = (\text{diss. } H_2O_2) = 15.9$  kcal [4], for the reaction




we obtain  $\Delta Z_O = -17$  kcal. The most probable values of the standard free energy of formation and hydration of  $H_2O_2$  are equal to -25.2 and -6.7 kcal [4, 5] respectively.

From this we get



Since  $\Delta Z_{\text{hydr}O_2} = 4$  kcal [6]. Using known values for the standard entropies of  $H_2$ ,  $HO_2$  and  $H_2O_2$  (31.2; 55.17, and 55.66 E.U. respectively [7]), the entropies of hydration of  $HO_2$  and  $H_2O_2$  (-22 and -24.8 E.U. [4,8,9]) and the entropy of acid dissociation of  $H_2O_2$  (-25.8 E.U. [4]), we find the  $\Delta H_O$  and  $\Delta S_O$  for reactions (1) and (2).


We now calculate the  $\Delta S_O$  for the acid dissociation of  $HO_2$ . From the entropic cycle



we find  $\Delta S_{\text{hydrO}_2^-} - \Delta S_{\text{dissH}_2\text{O}_2} = -12.2 \text{ E.U.}$  The entropies of hydration of polyatomic ions monatomically increase in proportion to the decrease of the radius of the ion in solution [10, 11]:

ANION	$\text{OH}^-$	$\text{CN}^-$	$\text{NO}_3^-$	$\text{ClO}_4^-$
$\Delta S_{\text{hydr}}$	-40.6	-19.9	-19.5	-16.3
Radius of ion				
in solution Å	1.53	1.89	1.92	2.36

It is reasonable to assume that the radius of the  $\text{O}_2^-$  ion-radical in solution is intermediate between the radii of  $\text{CN}^-$  and  $\text{NO}_3^-$ . This indicates that the most probable value for the entropy of hydration of  $\text{O}_2^-$  is equal to the mean value (-19.7 E.U.) of the entropies of hydration for these anions. We then obtain -7.5 E.U. for the most probable value of  $\Delta S_{\text{dissH}_2\text{O}_2}$ . The free energy of dissociation of  $\text{H}_2\text{O}_2$  is equal to 6.65 kcal<sup>2</sup> ( $p^{\text{K}}_{\text{H}_2\text{O}_2} = 4.88$  [11]). For the enthalpy of dissociation we get  $\Delta H = 4.4 \text{ kcal}$ . As a result we find the values for standard entropies and enthalpies of reactions (5) and (6) (see the Table). With the aid of the cycle




we find  $E_{\text{H-O}_2\text{H}} - H_{\text{hydrH}_2\text{O}_2} = 76.3 \text{ kcal}$ .

/2660

Number	Reaction	$\Delta H^\circ$ , kcal	$\Delta S^\circ$ , E. U.	$\Delta Z^\circ$ , kcal	$E^\circ$ , V
1	$\frac{1}{2}\text{H}_2 + \text{HO}_2(\text{aq}) \rightarrow \text{H}_2\text{O}_2(\text{aq})$	-37.9	-16.9	-32.9	1.43
2	$\frac{1}{2}\text{H}_2 + \text{HO}_2(\text{aq}) \rightarrow \text{HO}_2^- + \text{H}^+$	-29.7	-42.7	-17.0	0.74
3	$\frac{1}{2}\text{H}_2 + \text{O}_2(\text{g}) \rightarrow \text{HO}_2(\text{aq})$	-8.4	-31.4	1.0	-0.04
4	$\frac{1}{2}\text{H}_2 + \text{O}_2(\text{aq}) \rightarrow \text{HO}_2(\text{aq})$	-5.4	-8.2	-3.0	0.13
5	$\frac{1}{2}\text{H}_2 + \text{O}_2(\text{g}) \rightarrow \text{O}_2^- + \text{H}^+$	-4.0	-38.9	7.7	-0.33
6	$\frac{1}{2}\text{H}_2 + \text{O}_2(\text{aq}) \rightarrow \text{O}_2^- + \text{H}^+$	-1.0	-15.7	3.7	-0.16
7	$\text{HO}_2(\text{g}) \rightarrow \text{HO}_2(\text{aq})$	-12.4	-22.0	-5.8	—
8	$\text{O}_2^-(\text{g}) \rightarrow \text{O}_2^-(\text{aq})$	-84.0	-19.7	-78.0	—
9	$\text{HO}_2 \rightleftharpoons \text{H}^+ + \text{O}_2^-$	4.4	-7.5	6.65	—

Assuming  $E_{H-O_2H} = 88.6$  kcal [12], we get  $\Delta H_{hydrH_2O_2} = -12.4$  kcal. According to [12] the energy error for the breaking of OH bonds in peroxide amounts to  $\pm 2$  kcal. But if  $E_{H-O_2H}$  is in fact equal to 90.6 kcal, then for  $\Delta H_{hydrH_2O_2}$  we would get 14.3 kcal. This is why a greater heat of hydration for peroxide is not very likely. When  $E_{H-O_2H} = 86.6$  kcal, the value becomes  $\Delta H_{hydrH_2O_2} = 10.3$  kcal, which is approximately equal to the heat of evaporation of water and also scarcely probable. The most reasonable hypothesis is that the heat of hydration of  $H_2O_2$  falls between the heat of evaporation of water and the heat of hydration of peroxide. The value we found meets this condition. Obviously,  $E_{H-O_2H} = 86.6$  kcal is more exact, but the error for the values of  $E_{H-O_2H}$  and  $\Delta H_{hydrH_2O_2}$  did not exceed 1 kcal. Using  $\Delta S_{hydrH_2O_2} = -22$  E.U. we find that  $\Delta Z_{hydrH_2O_2} = -5.8$  kcal. By means of the cycle



it is possible to obtain a sufficiently accurate value for the free energy of hydration of  $O_2^-$ , which is equal to -78 kcal, from which we find  $\Delta H_{hydrO_2^-} = -84$  kcal.

As the next procedure, we conducted additional verification of the values found. According to [14], the ratio  $(k_{Fe^{3+} + O_2^-} / k_{dissH_2O_2}) / k_{Fe^{2+} + H_2O_2} = 7 \cdot 10^{-3}$  mol/l at 25°C, and the energy of activation for this ratio is equal to 1.2-4 kcal. We have  $k_{Fe^{3+} + O_2^-} = 1.1 \cdot 10^9$  /mol·sec. Since the limiting phase of the reaction  $Fe^{2+} + H_2O_2$  is the replacement of water in the hydrated shell of the  $Fe^{2+}$  ion, the energy of activation  $E_{Fe^{2+} + H_2O_2}$  must be close to the energy of activation for replacement, i.e., 8.3 kcal [15]. Using  $\Delta H_{dissH_2O_2} = 4.4$  kcal, we arrive at the result

$E_{\text{Fe}^{3+} + \text{O}_2^-} = 5.1-7.9 \text{ kcal}$ . Assuming  $E_{\text{Fe}^{3+} + \text{O}_2^-}$  to be  $= 6.5 \text{ kcal}$ , we obtain  $k_{\text{Fe}^{3+} + \text{O}_2^-} = 10^{14} e^{-6500/RT}$ . However, taking account of a considerable positive input into the energy of activation, which corresponds to the reaction between inversely charged molecules (in the given case from three to six orders of pre-exponential magnitude), we must recognize the value found of the energy of activation as completely reasonable. In [8] the value  $\Delta S_{\text{dissH}_2\text{O}_2} = -27 \text{ E.U.}$  was used, giving  $\Delta H_{\text{dissH}_2\text{O}_2} = -1.4 \text{ kcal}$  and  $E_{\text{Fe}^{3+} + \text{O}_2^-} = 10.9 - 13.7 \text{ kcal}$ , which signifies an abnormally high value for the pre-exponential factor and is, obviously, unreliable.

According to [16], the rate constant for the reaction  $\text{Cu}^{2+} + \text{O}_2^-$  is equal to  $2 \cdot 10^9 \text{ l/mol} \cdot \text{sec}$  at  $25^\circ\text{C}$ . According to our data, we find  $\Delta H = 2.7 \text{ kcal}$  for this reaction, which allows a possibility of a low energy of activation for it. For example, if the energy of activation of the reverse reaction is  $\sim 2 \text{ kcal}$ , then, for the reaction under consideration, we obtain  $E_{\text{act}} \sim 5 \text{ kcal}$ , which gives  $k_{\text{Cu}^{2+} + \text{O}_2^-} = 10^{13} e^{-5000/RT}$ . Because of reasons analogous to that shown above for the reaction  $\text{Fe}^{3+} + \text{O}_2^-$ , the value of the pre-exponential factor is also reliable here. But if the data in [8] is used, then for this reaction we obtain  $\Delta H = 8.5 \text{ kcal}$  and  $E_{\text{act}} \sim 10.5 \text{ kcal}$ , which, obviously, is not very likely for so rapid a reaction. The results of the calculation of the thermodynamic characteristics of  $\text{H}_2\text{O}_2$  in an aqueous solution are presented in the table. The error for the values of  $E_0$ , which correspond to reactions (1) - (6), apparently does not exceed  $0.01 \text{ V}$ . The error for the values of  $\Delta H_0$  and  $\Delta S_0$ , evidently, does not exceed  $1 \text{ kcal}$  and  $3-4 \text{ E.U.}$ , respectively. Approximately, then, this is the accuracy of the values corresponding to reactions (7) - (9).

Institute of Chemical Kinetics and Combustion  
USSR Academy of Sciences  
Siberian Division  
Novosibirsk

Received  
December 21, 1971



## REFERENCES

1. V.M. Berdnikov, O.S. Zhusavleva, Kinetika i Kataliz (Kinetics and Catalysis), in press.
2. G.G. Jayson, J.P. Keene, D.A. Stirling, A.J. Swallow, Trans, Faraday Soc., 65, 2454, 1969.
3. Yu.Yu. Lur'ye, Spravochnik po Analiticheskoy Khimii (Reference source on Analytic Chemistry), "Khimiya" Press, Moscow, 1967.
4. U. Shamb, Ch. Setterfield, R. Ventuors, Perehis' Vodoroda (Hydrogen Peroxide), Foreign Lit. Press, Moscow, 1958.
5. M.Kh. Karapet'yants, M.L. Karapet'yants, Osnovnyye Termodinamicheskie Konstanty Neorganicheskikh i organicheskikh Beshchest v (Basic Thermodynamic Constants of Inorganic and Organic Substances), "Khimiya" Press, Moscow, 1968.
6. R.A. Pierotti, J. Phys. Chem., 69, 281, 1965.
7. A.V. Gurvich, Termodinamicheskiye svoystva individual'nykh veshchestv T. I, (Thermodynamic properties of individual substances, Vol. I.) AN SSSR Press, Moscow, 1962.
8. Yu.I. Skurlatoo, Zh. Fiz. Khimii (Journal of Physical Chemistry), 44, 548, 1970.
9. V.M. Berdnikov, N.M. Bazhin, Zh. Fiz. Khimii ([as above]), 44, 712, 1970.
10. Kratkiy spravochnik fiziko-khimicheskikh velichin (Short listing of Physical-Chemical Constants), "Khimiya" Press, Moscow, 1967.
11. D. Behar, D.B.J. Czapski, J. Rabani, L.M. Dorfman, H.A. Schwarz, J. Phys. Chem., 74, 3209, 1970.
12. V.I. Vedeneyev, Energiya razryva svyazey (Energy of the Breaking of Bonds), Moscow, 1962.

13. V.E. Conway, J.O.M. Bockris, in the collection  
Nekotoryye problemy sovremennoy elektrokhimii  
(Some Problems of Modern Electrochemistry), Foreign  
Literature Press, Moscow, 1958.
14. W.O. Barb, J.H. Bakendale, P. George, K.R. Hargrave,  
Trans. Faraday Soc., 47, 464, 1951.
15. N. Sutin, Ann. Rev. Phys. Chem., 17, 119, 1966.
16. Yu.N. Kozlor, V.M. Berdnikov, Zh. Fiz. Khimii  
(Journal of Physical Chemistry), in press.